

Nuclear uncertainties in the NeNa-MgAl cycles and production of ^{22}Na and ^{26}Al during nova outbursts

Jordi José

Departament de Física i Enginyeria Nuclear (UPC), Av. Víctor Balaguer, s/n, E-08800

Vilanova i la Geltrú (Barcelona), SPAIN

Institut d'Estudis Espacials de Catalunya (IEEC), Edifici Nexus-201, C/ Gran Capità 2-4,

E-08034 Barcelona, SPAIN

Alain Coc

Centre de Spectrométrie Nucléaire et de Spectrométrie de Masse, IN2P3-CNRS, Bât. 104,

F-91405 Orsay Campus, FRANCE

and

Margarita Hernanz

Institut d'Estudis Espacials de Catalunya (IEEC), CSIC Research Unit, Edifici Nexus-201,

C/ Gran Capità 2-4, E-08034 Barcelona, SPAIN

Received _____; accepted _____

Submitted to: ApJ Version: February 1, 2008

ABSTRACT

Classical novae eject significant amounts of nuclear processed material into the interstellar medium. Among the isotopes synthesized during such explosions, two radioactive nuclei deserve a particular attention: ^{22}Na and ^{26}Al . In this paper, we investigate the nuclear paths leading to ^{22}Na and ^{26}Al production during nova outbursts by means of an implicit, hydrodynamic code that follows the course of the thermonuclear runaway from the onset of accretion up to the ejection stage. New evolutionary sequences of ONe novae have been computed, using updated nuclear reaction rates relevant to ^{22}Na and ^{26}Al production. Special attention is focused on the role played by nuclear uncertainties within the NeNa and MgAl cycles in the synthesis of such radioactive species. From the series of hydrodynamic models, which assume upper, recommended or lower estimates of the reaction rates, we derive limits on the production of both ^{22}Na and ^{26}Al . We outline a list of nuclear reactions which deserve new experimental investigations in order to reduce the wide dispersion introduced by nuclear uncertainties in the ^{22}Na and ^{26}Al yields.

Subject headings: novae, cataclysmic variables — nuclear reactions, nucleosynthesis, abundances

1. Introduction

The thermonuclear runaway model has been successful in reproducing the *gross* features of nova outbursts. According to this widely accepted scenario, classical novae are produced by thermonuclear runaways (hereafter, TNRs) that take place in the white dwarf component of a close binary system. The large, main sequence companion overfills its Roche lobe, providing matter outflows through the inner Lagrangian point that lead to the formation of an accretion disk around the white dwarf. A fraction of this H-rich matter lost by the companion ultimately ends up on top of the white dwarf, where it is gradually compressed as accretion goes on. The piling up of matter heats the envelope up to the point when ignition conditions to drive a TNR are reached.

An extended set of hydrodynamic computations of classical nova outbursts has been performed during the last 25 years (see Starrfield et al. 1972, for the first hydrodynamic study of the TNR model, and the recent papers by Kovetz & Prialnik 1997, Starrfield et al. 1998, José & Hernanz 1998, and references therein), for a wide range of white dwarf masses and initial luminosities, mass accretion rates and initial chemical compositions. From the nucleosynthesis viewpoint, these computations have been able to identify two types of nova outbursts, those occurring in CO or in ONe white dwarfs. The latter ones have provided a framework for the origin of the high concentrations of Ne and more massive isotopes found in the spectra of some well-observed novae, such as V693 CrA 1981, V1370 Aql 1982, QU Vul 1984 No. 2, V838 Her 1991 or V1974 Cyg 1992 (Livio & Truran 1994; Gehrz et al. 1998). Among the isotopes synthesized in these so-called Ne (or ONe) novae, two radioactive species have raised a particular astrophysical interest: ^{22}Na and ^{26}Al .

The potential role of ^{22}Na for diagnosis of nova outbursts was first suggested by Clayton & Hoyle (1974). It decays to a short-lived excited state of ^{22}Ne (with a lifetime of $\tau = 3.75$ yr), which de-excites to its ground state by emitting a γ -ray photon of 1.275 MeV.

Through this mechanism, nearby ONe novae within a few kiloparsecs from the Sun may provide detectable γ -ray fluxes. Several experimental verifications of this γ -ray emission at 1.275 MeV from nearby novae have been attempted in the last twenty years, using balloon-borne experiments (Leventhal et al. 1977), and detectors on-board satellites such as HEAO-3 (Mahoney et al. 1982), SMM (Leising et al. 1988), and CGRO (Leising 1993; Iyudin et al. 1995), from which upper limits on the ejected ^{22}Na have been derived. In particular, the observations performed with the COMPTEL experiment on-board CGRO of five recent Ne-type novae (Nova Her 1991, Nova Sgr 1991, Nova Sct 1991, Nova Pup 1991 and Nova Cyg 1992. Iyudin et al. 1995), as well as observations of *standard* CO novae, have led to an upper limit of $3.7 \times 10^{-8} M_{\odot}$ for the ^{22}Na mass ejected by any nova in the Galactic disk. This restrictive limit has posed some constraints on pre-existing theoretical models of classical nova explosions.

^{26}Al is another unstable nucleus, with a lifetime of $\tau = 1.04 \times 10^6$ years, that decays from ground state to the first excited level of ^{26}Mg , which in turn de-excites to its ground state by emitting a gamma-ray photon of 1.809 MeV. This characteristic gamma-ray signature, first detected in the Galactic Center by the HEAO-3 satellite (Mahoney et al. 1982, 1984), has been confirmed by other space missions like SMM (Share et al. 1985) and by several balloon-borne experiments. The most recent measurements made with COMPTEL have provided a map of the 1.809 MeV emission in the Galaxy (Diehl et al. 1995, 1997; Prantzos & Diehl 1996). The inferred $1 - 3 M_{\odot}$ of Galactic ^{26}Al are, according to the observed distribution, mainly attributed to young progenitors, such as massive AGB stars, type II supernovae and Wolf-Rayet stars. More recent analyses of the 1.809 MeV COMPTEL map reveal a correlation between this map and the COBE/DMR maps, tracing free-free emission, thus confirming that the main contributors to the Galactic ^{26}Al are massive stars (Knödlseider 1997). However, a potential contribution from novae or low-mass AGB stars cannot be ruled out (see José, Hernanz & Coc 1997, for a recent analysis of ^{26}Al

production in classical novae).

First estimates of the ^{22}Na and ^{26}Al production in novae were performed by different groups, using simplified one-zone models with representative temperature and density profiles. Hillebrandt & Thielemann (1982) and Wiescher et al. (1986) suggested already that classical novae might produce significant amounts of ^{26}Al , not enough to represent major Galactic sources, but relevant to account for the observed isotopic anomalies found in some meteorites. New parametrized calculations by Weiss & Truran (1990) and Nofar, Shaviv & Starrfield (1991), revealed that nova envelopes previously enhanced in heavy elements (from Ne to Mg) produce large amounts of ^{22}Na and ^{26}Al . Since this metal enrichment is expected to result from dredge-up of core material, Weiss & Truran suggested that massive ONeMg white dwarfs (the ones attaining the highest peak temperatures and, therefore, the most efficient dredge-up) are likely to provide the largest abundances of both ^{22}Na and ^{26}Al in the ejecta. Politano et al. (1995) revisited this scenario using hydrodynamic computations. They reported on a strong anticorrelation between ^{22}Na and ^{26}Al production: novae that produce the largest amounts of ^{22}Na (i.e., massive white dwarfs) are not the same as those accounting for the largest yields of ^{26}Al (i.e., low-mass white dwarfs). The obtained ^{22}Na yields range between 5×10^{-5} and 5×10^{-3} , by mass. Assuming that the whole envelope ($\sim 10^{-4} - 10^{-5} M_{\odot}$) is ejected during the outburst, they concluded that nearby ONeMg novae with $M_{wd} \geq 1.25 M_{\odot}$ should produce detectable ^{22}Na γ -rays for CGRO, a prediction not confirmed so far (Iyudin et al. 1995). Their results showed also a significant production of ^{26}Al (i.e., $(19.6 - 7.5) \times 10^{-3}$, by mass, corresponding to ONeMg white dwarfs with masses between $1.0 - 1.35 M_{\odot}$, respectively), which could account for a major fraction of the Galactic ^{26}Al .

Recent hydrodynamic computations of ONe novae (José, Hernanz & Coc 1997; José & Hernanz 1997, 1998) using updated initial compositions and nuclear reaction rates, have

led to a significant reduction of both ^{26}Al and ^{22}Na ejected during nova outbursts. In particular, a mean mass fraction of 1×10^{-4} of ^{22}Na is found in the $1.25 M_{\odot}$ ONe Model (with $M_{\text{ejec}}(^{22}\text{Na}) = 1.3 \times 10^{-9} M_{\odot}$), whereas a maximum value of 6×10^{-4} results from the $1.35 M_{\odot}$ ONe Model ($M_{\text{ejec}}(^{22}\text{Na}) = 2.6 \times 10^{-9} M_{\odot}$). The corresponding peak fluxes in the 1.275 MeV ^{22}Na line, below 10^{-5} photons $\text{s}^{-1} \text{cm}^{-2}$ for novae at 1 kpc, turn out to be too low to be detected with OSSE or COMPTEL but represent potential targets for the nearby future INTEGRAL mission (Hernanz et al. 1997; Gómez-Gomar et al. 1998). Concerning ^{26}Al , yields ranging from 2×10^{-3} to 2×10^{-4} by mass have been obtained in a series of ONe nova models with masses between $1.15 - 1.35 M_{\odot}$. Contribution of novae to the Galactic ^{26}Al is limited to $0.1 - 0.4 M_{\odot}$ (José, Hernanz & Coc 1997). Nevertheless, a larger contribution cannot be ruled out if the (uncertain) lower limit for ONe white dwarfs is reduced down to $1.0 M_{\odot}$ (José & Hernanz 1998).

Other hydrodynamic computations performed by Starrfield et al. (1997, 1998), using also updated nuclear reaction rates and opacities, have modified their previous estimates (Politano et al. 1995). The expected abundance of ^{22}Na in the ejecta has risen up to $(2 - 3) \times 10^{-3}$, by mass, when $1.25 M_{\odot}$ ONeMg white dwarfs are adopted, high enough to be detected by CGRO, provided that all the accreted envelope ($3 \times 10^{-5} M_{\odot}$) is ejected. On the other hand, the improved input physics translates into a factor of 10 reduction on the synthesis of ^{26}Al , in better agreement with the analysis of the 1.809 MeV emission map provided by COMPTEL, and also with the results previously reported by José, Hernanz & Coc (1997).

Whereas the agreement between the different groups on the expected contribution of classical novae to the Galactic ^{26}Al has significantly increased, there remains some discrepancy concerning the amount of ^{22}Na present in the ejecta. Since the synthesis of both ^{22}Na and ^{26}Al is very dependent on the adopted nuclear reaction rates, the large

uncertainties present in some key reactions of both NeNa and MgAl cycles (Coc et al. 1995; Prantzos & Diehl 1996), may significantly modify the expected yields. In particular, the study of the influence of a given reaction rate on ^{22}Na production is not trivial due to the two possible modes of formation and their interplay with convection (Coc et al. 1995). Moreover, during a nova outburst, thermodynamic conditions change on a short timescale so that nuclear reactions are never close to equilibrium. Due to obvious experimental difficulties, reaction rates involving short-lived radioactive nuclei of the NeNa-MgAl group are in general poorly known and the associated uncertainties may reach many orders of magnitude. It is thus important to know how the yields of important isotopes like ^{22}Na and ^{26}Al are affected by those uncertainties. In this paper, series of hydrodynamic nova models have been computed assuming upper, recommended and lower estimates of the reaction rates, from which limits on the production of both ^{22}Na and ^{26}Al are derived. The present analysis is focused on capture rates on radioactive nuclei, and for a temperature domain in the range $T_8 \simeq 0.5 - 3.5$. We refer to the NACRE compilation (Angulo et al. 1998) for a more general discussion concerning capture rates on stable isotopes, and a wider temperature range. For convenience, we use the terms NeNa cycle and MgAl cycle to denote the reactions involved in ^{22}Na and ^{26}Al formation. However, because of the high $^{23}\text{Na}(p,\gamma)$ and $^{27}\text{Al}(p,\gamma)$ rates, they cannot be considered as genuine *cycles*.

In Section 2, we outline some details of the method of computation and input physics. A detailed analysis of the synthesis of ^{22}Na and ^{26}Al in classical novae, together with the study of the role played by specific nuclear reactions of the NeNa-MgAl cycles, is given in Sections 3 & 4. Constraints on the production of ^{22}Na and ^{26}Al assuming lower, recommended and upper rates for some key reactions are derived in Section 5. The most relevant conclusions of this paper are summarized in Section 6. A detailed Appendix focused on the nuclear physics aspects of the reaction rates within the NeNa and MgAl cycles follows.

2. Model and Input physics

Evolutionary sequences of nova outbursts have been calculated by means of an updated version of the code SHIVA (see José 1996; José & Hernanz 1998), a one-dimensional, implicit, hydrodynamical code in lagrangian formulation, that follows the course of the outburst from the onset of accretion up to the expansion and ejection stages. The code solves the standard set of differential equations for hydrodynamical evolution: conservation of mass, momentum and energy, energy transport by radiation and convection, plus the definition of the lagrangian velocity, including a time-dependent formalism for convective transport whenever the characteristic convective timescale becomes larger than the integration time step (Wood 1974). Partial mixing between adjacent convective shells is treated by means of a diffusion equation (see Prialnik, Shara & Shaviv 1979, for the formalism). The code is linked to a reaction network, which follows the detailed evolution of 100 nuclei, ranging from ^1H to ^{40}Ca , through 370 nuclear reactions, with updated rates, and screening factors from Graboske et al. (1973) and DeWitt, Graboske & Cooper (1973). As suggested by Politano et al. (1995), the matter transferred from the companion is assumed to be solar-like, and is mixed in a given fraction with the outermost shells of the underlying core by means of an unknown mechanism (either shear mixing, diffusion or a convective multidimensional process). The composition of the underlying core has been taken from recent detailed evolutionary models in the case of ONe white dwarfs, which are the main contributors to ^{22}Na and ^{26}Al synthesis. These stars are made basically of ^{16}O and ^{20}Ne (Domínguez, Tornambé & Isern 1993; Ritossa, García-Berro & Iben 1996), with smaller traces of ^{23}Na , $^{24,25}\text{Mg}$, ^{27}Al and other species. This issue plays a crucial role in the resulting nucleosynthesis, and should be taken into account in order to compare results obtained by different groups. In particular, the ONeMg models computed by Starrfield et al. (1998) have an initial composition of the white dwarf core based on old nucleosynthesis calculations of C-burning from Arnett & Truran (1969), which is richer in ^{24}Mg and ^{20}Ne

than the one adopted in this paper (see Table 1).

3. Synthesis of ^{22}Na in Classical Novae

The high temperatures attained during ONe nova outbursts (with peak values within $2 - 3.5 \times 10^8$ K), allow a noticeable nuclear activity in the NeNa and MgAl cycles, which results on a significant production of species of astrophysical interest, such as ^{22}Na and ^{26}Al . In this Section, we will describe the main mechanisms of ^{22}Na synthesis, through a detailed analysis of a $1.25 M_{\odot}$, ONe white dwarf, which accretes solar-like matter at a rate $\dot{M} = 2 \times 10^{-10} M_{\odot}.\text{yr}^{-1}$, assuming a 50% degree of mixing with the ONe core (Model ONe5, in José & Hernanz 1998). Snapshots of the evolution of several isotopes relevant to ^{22}Na synthesis (i.e., $^{20,21,22}\text{Ne}$, $^{21,22,23}\text{Na}$ and $^{22,23}\text{Mg}$) are shown in Figure 1.

3.1. Main nuclear reactions involved in ^{22}Na production

At the onset of accretion, the evolution of ^{22}Na is mainly dominated by the chain of nuclear reactions $^{20}\text{Ne}(p, \gamma)^{21}\text{Na}(\beta^+)^{21}\text{Ne}(p, \gamma)^{22}\text{Na}(\beta^+)^{22}\text{Ne}(p, \gamma)^{23}\text{Na}$ (i.e., the cold mode of the NeNa cycle. See Fig. 2). When the temperature at the burning shell reaches $T_{bs} = 5 \times 10^7$ K (Fig. 1, panel 1), the main nuclear reaction of the NeNa cycle is $^{21}\text{Ne}(p, \gamma)^{22}\text{Na}$, which significantly reduces the amount of ^{21}Ne . In fact, when $T_{bs} = 7 \times 10^7$ K (panel 2), the amount of ^{21}Ne is already too small to maintain the main mechanism for ^{22}Na synthesis. Therefore, ^{22}Na will begin to decrease near the burning shell due to proton captures, following the rise of temperature toward peak value (panels 3 & 4).

At $T_{bs} = 1 \times 10^8$ K (panel 3), the amount of ^{21}Ne has already decreased below 10^{-6} by mass, except at the outer envelope, where some ^{21}Ne is synthesized from the β^+ -decay of ^{21}Na (previously built up as $^{20}\text{Ne}(p, \gamma)^{21}\text{Na}$). ^{21}Na and ^{23}Mg increase due to $^{20}\text{Ne}(p,$

γ) ^{21}Na and $^{22}\text{Na}(\text{p}, \gamma)^{23}\text{Mg}$, respectively. Destruction of ^{22}Na through (p, γ) reactions goes on but, due to convection, ^{22}Na shows a nearly flat profile throughout the envelope.

When $T_{bs} = 2 \times 10^8$ K (panel 4), there is a dramatic decline in ^{23}Na (due to (p, γ) and (p, α) reactions) and in the ^{22}Na abundance (by $^{22}\text{Na}(\text{p}, \gamma)^{23}\text{Mg}$). ^{22}Ne has slightly decreased. Also noticeable is the increase of ^{21}Na (since $^{20}\text{Ne}(\text{p}, \gamma)^{21}\text{Na}$ dominates destruction from both $^{21}\text{Na}(\beta^+)^{21}\text{Ne}$ and $^{21}\text{Na}(\text{p}, \gamma)^{22}\text{Mg}$), which plays a crucial role in the synthesis of ^{22}Na at the late stages of the outburst. Both $^{22,23}\text{Mg}$ increase due to proton captures on $^{21,22}\text{Na}$ respectively.

33 seconds later, the temperature at the burning shell attains its peak value, $T_{bs, max} = 2.44 \times 10^8$ K (panel 5). The amount of ^{21}Na is maintained by a quasi-equilibrium between $^{21}\text{Na}(\text{p}, \gamma)^{22}\text{Mg}$ and $^{20}\text{Ne}(\text{p}, \gamma)^{21}\text{Na}$. The mean amount of ^{22}Na increases due to $^{22}\text{Mg}(\beta^+)^{22}\text{Na}$, previously transported by convection to the outer, cooler layers of the envelope. The amount of ^{23}Na decreases due to both (p, γ) and (p, α) reactions, which dominate $^{22}\text{Ne}(\text{p}, \gamma)^{23}\text{Na}$ as well as $^{23}\text{Mg}(\beta^+)^{23}\text{Na}$. Since the peak temperature achieved in the burning shell is not extremely high, ^{20}Ne remains nearly unchanged (the typical temperature for ^{20}Ne burning exceeds $\sim 4 \times 10^8$ K). With respect to the other neon isotopes, ^{21}Ne increases (mainly due to the β^+ -decay of ^{21}Na at the outer shells), whereas ^{22}Ne is destroyed by $^{22}\text{Ne}(\text{p}, \gamma)^{23}\text{Na}$. The two magnesium isotopes $^{22,23}\text{Mg}$ increase as a result of proton captures onto $^{21,22}\text{Na}$.

Shortly after, due to the sudden release of energy from the short-lived β^+ -unstable nuclei ^{13}N , $^{14,15}\text{O}$, and ^{17}F , the envelope begins to expand. The role played by (p, γ) and (p, α) reactions is therefore reduced following the drop in temperature, whereas β^+ -decays progressively dominate the evolution. The abundances of ^{21}Na and $^{22,23}\text{Mg}$ decrease as a result of such β^+ -decays, which in turn increase the amount of ^{21}Ne and $^{22,23}\text{Na}$ (panels 6 & 7).

At the final stages of the outburst (panel 8), as the envelope expands and cools down, most of the remaining nuclear activity in the NeNa cycle is due to β^+ -decays, such as $^{21}\text{Na}(\beta^+)^{21}\text{Ne}$, $^{22}\text{Mg}(\beta^+)^{22}\text{Na}$ or $^{23}\text{Mg}(\beta^+)^{23}\text{Na}$. The resulting mean abundance of ^{22}Na in the ejected shells of this $1.25 M_{\odot}$ ONe Model is $X(^{22}\text{Na}) = 9.6 \times 10^{-5}$ by mass, which corresponds to $1.3 \times 10^{-9} M_{\odot}$ of ^{22}Na ejected into the interstellar medium. Other species of the NeNa cycle present in the ejecta are ^{20}Ne ($X(^{20}\text{Ne}) = 0.18$, a mass fraction slightly higher than its initial value because of the operation of the $^{23}\text{Na}(p, \alpha)^{20}\text{Ne}$ reaction. See Table 1), $^{21,22}\text{Ne}$ ($X(^{21}\text{Ne}) = 3.5 \times 10^{-5}$, and $X(^{22}\text{Ne}) = 1.0 \times 10^{-3}$), and ^{23}Na (with $X(^{23}\text{Na}) = 1.4 \times 10^{-3}$).

3.2. Effect of the reaction rates on the synthesis of ^{22}Na

From the abovementioned nuclear physics viewpoint, the synthesis of ^{22}Na is mainly controlled by four reactions: $^{20}\text{Ne}(p, \gamma)^{21}\text{Na}$, $^{21}\text{Na}(p, \gamma)^{22}\text{Mg}$, $^{21}\text{Ne}(p, \gamma)^{22}\text{Na}$ and $^{22}\text{Na}(p, \gamma)^{23}\text{Mg}$ (Fig. 2), for which recent updates to the reaction rates are available. Despite energy production during nova outbursts is not very dependent on the specific prescriptions adopted for such rates (see Section 5.1), they play a crucial role on the accompanying nucleosynthesis, since changes in the reaction paths are expected. Several test models have been computed to analyse the role played by different nuclear reactions on the synthesis of ^{22}Na (as well as ^{26}Al). Main results concerning ^{22}Na and ^{26}Al production, compared with the abundances found in José & Hernanz (1998) with previous prescriptions, are summarized in Table 2.

$^{20}\text{Ne}(p, \gamma)^{21}\text{Na}$ is the slowest proton capture reaction on any stable neon and sodium isotope. According to the adopted composition for the ONe core (Ritossa, García-Berro & Iben 1996), ^{20}Ne is the most abundant Ne-Na isotope. Therefore, $^{20}\text{Ne}(p, \gamma)^{21}\text{Na}$ limits ^{22}Na production in classical novae. The adopted $^{20}\text{Ne}(p, \gamma)^{21}\text{Na}$ rate is the Caughlan &

Fowler (1988) one, which is based on Rolfs et al. (1975) data. In the considered domain of temperature, it is well known. The estimated uncertainty is of a factor of $\simeq 3$ (defined as the ratio between high and low estimates within the domain of temperature considered), according to the new compilation of nuclear reaction rates (Angulo et al. 1998), but more likely a factor of $\simeq 1.5$ only, as it is derived from short range standard extrapolations of experimental data (Rolfs et al. 1975). Therefore, nuclear uncertainties associated with $^{20}\text{Ne}(\text{p}, \gamma)^{21}\text{Na}$ should play no significant role on ^{22}Na production.

Concerning the $^{21}\text{Ne}(\text{p}, \gamma)^{22}\text{Na}$ rate, no significant effect of the reduction of the contribution of the $E_x = 6.384$ MeV level (Görres et al. 1983) with respect to Caughlan & Fowler (1988) has been found on ^{22}Na production (tested with a $1.25 M_\odot$ ONe white dwarf model). According to Angulo et al. (1998), other nuclear uncertainties scarcely affect the $^{21}\text{Ne}(\text{p}, \gamma)^{22}\text{Na}$ rate at moderate temperatures. On the contrary, much more uncertain is the $^{22}\text{Ne}(\text{p}, \gamma)^{23}\text{Na}$ rate, poorly known in the range of temperatures of interest for nova outbursts (Angulo et al. 1998). Nevertheless, due to the low initial ^{22}Ne abundance with respect to ^{23}Na , and also to the negligible $^{22}\text{Na}(\beta^+)^{22}\text{Ne}$ decay ($\tau = 3.75$ yr) during the TNR, the nuclear uncertainties affecting the $^{22}\text{Ne}(\text{p}, \gamma)^{23}\text{Na}$ rate are not relevant for ^{22}Na production in novae.

Recent experimental investigations of the $^{22}\text{Na}(\text{p}, \gamma)^{23}\text{Mg}$ reaction (Seuthe et al. 1990; Schmidt et al. 1995; Stegmüller et al. 1996) have provided a firmer basis for the determination of its rate. Since the new rate is lower than the old Caughlan & Fowler’s (1988) analytic fit (i.e., one order of magnitude for $T_8 > 1$), destruction of ^{22}Na by means of (p, γ) reactions is reduced. In a test model, consisting of a $1.25 M_\odot$ ONe white dwarf (Model ONe5, José & Hernanz 1998), the mean abundance of ^{22}Na in the ejecta increases by a factor of ~ 3 when the new $^{22}\text{Na}(\text{p}, \gamma)^{23}\text{Mg}$ rate (Stegmüller et al. 1996), instead of the CF88 one, is adopted (see Table 2). It is worth noticing that significant nuclear

uncertainties affect this rate (a factor ranging from 3 to 6 for $T_8 > 1$), which turn out to be crucial in order to derive ranges of ^{22}Na production during nova outbursts. According to Stegmüller et al. (1996), this uncertainty is mainly due to the possible effect of a resonance at $E_p = 225$ keV. Below $T_8=1$, the uncertainty reaches three orders of magnitude but the rate remains small enough to prevent destruction of ^{22}Na .

Three test models have been computed to analyze the role played by $^{21}\text{Na}(p, \gamma)^{22}\text{Mg}$, in view of the nuclear uncertainties present in this rate, in particular the estimated strength of the first $E_x = 5.714$ MeV level above the proton threshold (see Appendix, section A.1). Calculations assume ONe white dwarfs of masses 1.15, 1.25 and 1.35 $M_\odot$, and the same input physics than Models ONe3, ONe5 and ONe6 described in José & Hernanz (1998), but reducing the $^{21}\text{Na}(p, \gamma)^{22}\text{Mg}$ rate given by CF88 by a factor of 100 (similar, for novae, to the lower rate given in the Appendix). It results in a significant increase in the ^{22}Na abundances present in the ejecta (a factor of ~ 2 to 3, in the 1.15 and 1.25 $M_\odot$ Models, and a factor of ~ 1.2 , in the 1.35 $M_\odot$ Model. Table 2), as compared with the values found with the standard CF88 rate. This effect can be interpreted as follows: when the $^{21}\text{Na}(p, \gamma)^{22}\text{Mg}(\beta^+)^{22}\text{Na}$ rate is reduced by a factor of 100, the alternative path, $^{21}\text{Na}(\beta^+)^{21}\text{Ne}(p, \gamma)^{22}\text{Na}$, is favored. In this case, ^{22}Na production is delayed to a time when the envelope is already expanding and cooling down (contrary to the case when the higher $^{21}\text{Na}(p, \gamma)^{22}\text{Mg}$ rate is adopted). As a result, a major fraction of ^{22}Na survives. This, in turn, explains the lower effect found in the 1.35 $M_\odot$ Model, caused by the higher temperatures achieved in the envelope (with $T_{peak} = 3.2 \times 10^8$ K) which remain high enough at the time when ^{22}Na is synthesized. One should note that this effect (increase in the ^{22}Na yield when the $^{21}\text{Na}(p, \gamma)^{22}\text{Mg}$ rate is reduced) was not foreseen and stresses the importance of full hydrodynamical calculations.

Other reactions that may be involved in the synthesis of ^{22}Na (and ^{23}Na) are $^{22}\text{Mg}(p, \gamma)^{23}\text{Al}$, $^{23}\text{Al}(p, \gamma)^{24}\text{Si}$, or $^{23}\text{Mg}(p, \gamma)^{24}\text{Al}$. However, no significant effect on the ^{22}Na

production is found when using updated rates for such reactions (see Appendix and Table 2), because of the limited nuclear flow they conduct for nova conditions.

According to this analysis, full evolutionary sequences of nova outbursts, from the onset of accretion up to the ejection stage, have been performed taking into account the low, recommended and high estimates to the $^{21}\text{Na}(\text{p}, \gamma)^{22}\text{Mg}$, $^{22}\text{Na}(\text{p}, \gamma)^{23}\text{Mg}$ and $^{21}\text{Ne}(\text{p}, \gamma)^{22}\text{Na}$ rates (as well as the new $^{23}\text{Mg}(\text{p}, \gamma)^{24}\text{Al}$ rate derived by Kubono, Kajino & Kato 1995), in order to derive a range of ^{22}Na yields, resulting purely from nuclear physics uncertainties. Results are summarized in Section 5.

4. Synthesis of ^{26}Al in Classical Novae

In this Section, we will analyze the nuclear paths leading to ^{26}Al synthesis. As for ^{22}Na production, we will describe the course of the $1.25 M_{\odot}$, ONe nova model (José & Hernanz 1998, Model ONe5). Snapshots of the evolution of several isotopes relevant to ^{26}Al synthesis (i.e., $^{24,25,26}\text{Mg}$, $^{25,27}\text{Al}$, $^{26,27}\text{Si}$ and the ground and isomeric states for ^{26}Al , hereafter $^{26}\text{Al}^g$ and $^{26}\text{Al}^m$) are shown in Figure 3.

4.1. Main nuclear reactions involved in ^{26}Al production

Nucleosynthesis of ^{26}Al is complicated by the presence of a short lived ($\tau_{1/2} = 6.3 \text{ s}$) spin isomer. At low temperatures ($T_8 \lesssim 4$), the ^{26}Al ground and isomeric states do not reach thermal equilibrium and must be treated as two separate isotopes (Ward & Fowler 1980, see also Coc & Porquet 1998).

The nuclear activity in the MgAl cycle at the early phases of the TNR, when the temperature at the burning shell is $T_{bs} = 5 \times 10^7 \text{ K}$, is dominated by $^{25}\text{Mg}(\text{p}, \gamma)$, which

leads to both ^{26}Al ground and isomeric states. A significant amount of $^{26}\text{Al}^g$ is already synthesized at such temperatures (Figure 3, panel 1). Another aluminum isotope, ^{27}Al , is slightly enhanced with respect to its initial abundance by means of $^{26}\text{Al}^m(\beta^+)^{26}\text{Mg}(\text{p},\gamma)^{27}\text{Al}$. A similar trend is found when $T_{bs} = 7 \times 10^7$ K (panel 2): both $^{26}\text{Al}^g$ and ^{27}Al continue to rise (specially the first one, which increases by nearly a factor of 10 from the abundance shown in panel 1).

At $T_{bs} = 10^8$ K (panel 3), the evolution in the MgAl cycle is dominated by $^{24}\text{Mg}(\text{p},\gamma)^{25}\text{Al}(\beta^+)^{25}\text{Mg}$, which in turn accounts for a noticeable production of the β^+ -unstable nuclei ^{25}Al , and also for the increase in the mean $^{26}\text{Al}^g$ abundance (by means of $^{25}\text{Mg}(\text{p},\gamma)^{26}\text{Al}^{g,m}$). A nearly flat profile of $^{26}\text{Al}^g$ results from convective mixing, which extends already throughout the whole envelope.

A major change in the dominant nuclear path is found when $T_{bs} = 2 \times 10^8$ K (panel 4). The temperature attained near the burning shell is high enough to allow (p, γ) reactions to proceed efficiently. In particular, the abundance of ^{24}Mg is reduced by a factor of ~ 1000 . The isomer $^{26}\text{Al}^m$ exceeds already 10^{-3} by mass for most of the envelope. A significant fraction of ^{25}Al is transformed through proton captures into ^{26}Si , which will increase the abundance of $^{26}\text{Al}^m$, and in turn that of ^{27}Al in the late phases of the TNR. The final $^{27}\text{Al}/^{26}\text{Al}^g$ ratio will reflect a competition between two different paths: ^{24}Mg (fed by $^{23}\text{Na}(\text{p},\gamma)^{24}\text{Mg}$) is transformed by proton captures into ^{25}Al , which can either decay into ^{25}Mg or capture another proton, leading to ^{26}Si . Only the first channel accounts for $^{26}\text{Al}^g$ synthesis, whereas ^{27}Al can be produced by both paths, following $^{25}\text{Mg}(\text{p},\gamma)^{26}\text{Al}^{g,m}(\text{p},\gamma)^{27}\text{Si}(\beta^+)^{27}\text{Al}$, or $^{26}\text{Si}(\beta^+)^{26}\text{Al}^m(\beta^+)^{26}\text{Mg}(\text{p},\gamma)^{27}\text{Al}$. Anyway, because of the large abundance of ^{25}Mg in the envelope at this stage, proton captures on the seed ^{25}Mg become the major source of $^{26}\text{Al}^g$. Some leakage from the MgAl cycle due to $^{26}\text{Al}^{g,m}(\text{p},\gamma)^{27}\text{Si}$ is also obtained at this stage of the outburst.

When the burning shell attains its peak temperature ($T_{bs,max} = 2.44 \times 10^8$ K. Panel 5), most of the MgAl isotopes show a significant reduction near the burning shell because of (p, γ) reactions. The dominant paths at this stage are $^{25}\text{Mg}(\text{p},\gamma)^{26}\text{Al}$, $^{27}\text{Al}(\text{p},\gamma)^{28}\text{Si}$ and $^{26,27}\text{Si}(\text{p},\gamma)$ are much faster than the corresponding β^+ -decays). Following the course of the outburst (panels 6 to 8), as the envelope expands and cools down, proton capture reactions are progressively reduced. Therefore, the late time evolution is mainly dominated by β^+ -decays, such as $^{26}\text{Si}(\beta^+)^{26}\text{Al}^m$, $^{27}\text{Si}(\beta^+)^{27}\text{Al}$, $^{25}\text{Al}(\beta^+)^{25}\text{Mg}$ and $^{26}\text{Al}^m(\beta^+)^{26}\text{Mg}$, which in turn, increase the final amounts of $^{25,26}\text{Mg}$ and ^{27}Al . The mean abundance of ^{26}Al in the ejecta of this 1.25 $M_\odot$, ONe white dwarf model is $X(^{26}\text{Al}) = 5.4 \times 10^{-4}$ by mass, which translates into $7.6 \times 10^{-9} M_\odot$ of ^{26}Al ejected into the interstellar medium. Other isotopes of the MgAl group present in the ejecta are ^{27}Al ($X(^{27}\text{Al}) = 2 \times 10^{-3}$, half the initial abundance. See Table 1), ^{25}Mg (2.4×10^{-3}), ^{26}Mg (2.8×10^{-4}), and ^{24}Mg (2×10^{-4}).

4.2. Effect of the reaction rates on the synthesis of ^{26}Al

The previous analysis has revealed that several isotopes should be considered as potential seeds for ^{26}Al synthesis: ^{24}Mg , ^{25}Mg and, to some extent, ^{23}Na and ^{22}Ne . In this case, the number of nuclear reactions involved in the synthesis of ^{26}Al is rather large (see Fig. 2). Several test models of nova outbursts have been computed to analyze the role played by each one of the most relevant reactions. Main results are summarized in Table 2.

The $^{23}\text{Na}(\text{p},\gamma)^{24}\text{Mg}$ rate (in contrast with $^{23}\text{Na}(\text{p},\alpha)^{20}\text{Ne}$) has been strongly modified by the introduction of the upper limit for the $E_r^{cm}=0.138$ MeV resonance strength obtained by Görres, Wiescher & Rolfs (1989). When taking this upper limit, the rate is significantly increased up to $T_8 \simeq 2$ (Görres et al. 1989). Its effect has been tested by recomputing a 1.15 $M_\odot$, ONe white dwarf Model (i.e., Model ONe3. José & Hernanz 1998): an increase by

a factor of ~ 3 of the final ^{24}Mg yields results, which in turn leads to slightly larger mass fractions of both ^{25}Mg (factor 1.4) and ^{26}Al (factor 1.3). Some ^{27}Al is also overproduced (by a factor of 1.4) when this resonance is taken into account. Those effects are expected to be stronger for more massive white dwarfs.

At the temperatures attained in nova outbursts, $^{24}\text{Mg}(p,\gamma)^{25}\text{Al}$ may become even faster than $^{12}\text{C}(p,\gamma)^{13}\text{N}$. The $^{24}\text{Mg}(p,\gamma)^{25}\text{Al}$ rate is dominated by the presence of the $E_r^{cm} = 0.214$ MeV, $J^\pi = 1/2^+$ resonance and suffers little uncertainty.

$^{25}\text{Mg}(p,\gamma)^{26}\text{Al}^g$ is also of great importance since it is the only channel that leads to ^{26}Al in its ground state. One may expect that its potential uncertainties are directly reflected in the $^{26}\text{Al}^g$ yields. The new rate is significantly smaller by a factor of $\simeq 7$ (Iliadis et al. (1996)) than the Caughlan & Fowler 1988 one, below $T_8 \simeq 1$, but this has no consequences in the nova domain. Iliadis et al. (1996) reanalyzed available transfer reaction data concerning the lower lying resonances and concluded that the associated uncertainties are relatively small (i.e., a factor of 1.5–2). For $T_8 > 1.5$, available direct measurements lead also to very small uncertainties (a factor of 1.5, according to Iliadis et al. (1996)) and should not influence significantly ^{26}Al production. Moreover, since $^{25}\text{Mg}(p,\gamma)$ can lead either to $^{26}\text{Al}^g$ or $^{26}\text{Al}^m$ with a known branching ratio (Endt & Rolfs 1987, Iliadis et al. 1996), the $^{25}\text{Mg}(p,\gamma)^{26}\text{Al}^m$ rate is known with a similar precision as the $^{25}\text{Mg}(p,\gamma)^{26}\text{Al}^g$ one.

In nova nucleosynthesis, the most important resonance in $^{26}\text{Al}^g(p,\gamma)^{27}\text{Si}$ is the $E_r^{cm} = 0.188$ MeV, while uncertainties on the lower lying resonances have no influence (Coc et al. 1995). In our calculations, we use the only single direct measurement available (Vogelaar 1989) for this crucial resonance strength. Recently, the results of the proton transfer reaction that motivated this subsequent direct measurement have been published (Vogelaar et al. 1996) but the resonance strength deduced from this indirect measurement is strongly dependent on the adopted transferred angular momentum. Accordingly, we have checked

the influence of such resonance on the resulting ^{26}Al yields by reducing its strength by a factor of $1/3$, well within the range of values adopted by Angulo et al. (1998). We obtain a significant increase by a factor of ~ 2 in the final amount of ^{26}Al (for a $1.15 M_{\odot}$ ONe white dwarf Model. Table 2), which strongly stresses the need of additional direct measurements to confirm the results found by Vogelaar (1989).

$^{25}\text{Al}(p,\gamma)^{26}\text{Si}$ plays an important role on ^{26}Al synthesis, since it leads to the formation of the short-lived isomer (through $^{26}\text{Si}(\beta^+)^{26}\text{Al}^m$), instead of the long-lived ground state. We have checked the role played by the $^{25}\text{Al}(p,\gamma)^{26}\text{Si}$ rate, adopting the upper and lower estimates (see Appendix) provided by Coc et al. (1995). Results are compared with those obtained with the $^{25}\text{Al}(p,\gamma)^{26}\text{Si}$ rate given by Wiescher et al. (1986). Whereas the final amounts of ^{26}Al and ^{27}Al are very slightly enhanced when the lower rate is adopted (i.e., case A. Coc et al. 1995), a significant reduction of ^{26}Al by a factor of ~ 2 is obtained for the upper one (case C). In the first case, $^{25}\text{Al}(\beta^+)^{25}\text{Mg}$ becomes faster than $^{25}\text{Al}(p,\gamma)^{26}\text{Si}$ and, therefore, the final amount of ^{25}Mg (and in turn ^{26}Al) increases (Table 2).

Large uncertainties affect the $^{26}\text{Al}^m(p,\gamma)^{27}\text{Si}$ rate. The existing Caughlan & Fowler’s (1988) prescription for this rate results from a Hauser-Feshbach calculation, a statistical approach that is not reliable at low temperatures, where the contribution of isolated resonances dominates. Hence, the rate may differ by several orders of magnitude from the theoretical one at the temperatures achieved during nova outbursts. However, no noticeable effect on ^{26}Al production is found when the $^{26}\text{Al}^m(p,\gamma)^{27}\text{Si}$ rate, as given by CF88, is multiplied arbitrarily by a factor of 100, with the exception of a net reduction (factor ~ 2) on the final amount of ^{26}Mg .

Nuclear uncertainties significantly affect the $^{26}\text{Mg}(p,\gamma)^{27}\text{Al}$ rate. Since the initial amount of ^{26}Mg is of the same order of magnitude than that of ^{27}Al , and it is also fed continuously by $^{26}\text{Al}^m(\beta^+)^{26}\text{Mg}$ ($\tau = 9.15$ s), these uncertainties cannot be ignored as for

$^{22}\text{Ne}(\text{p},\gamma)^{23}\text{Na}$ (see Section 3.2). They are specially important around $T_8 = 0.5$, where the rate remains uncertain by a factor of $\simeq 100$ (Champagne et al. 1990), due to the unknown strength of an hypothetical resonance at $E_R^{cm} = 90$ keV. No influence on the final $^{26,27}\text{Al}$ yields results from the inclusion of the $^{26}\text{Mg}(\text{p},\gamma)^{27}\text{Al}$ recommended rate given by Champagne et al. (1990) (tested with a $1.15 M_\odot$ ONe white dwarf Model).

With respect to CF88, the $^{27}\text{Al}(\text{p},\gamma)^{28}\text{Si}$ rate has not changed: above $T_8=1$, uncertainties are limited to a factor of $\simeq 2$ (Angulo et al. 1998). On the contrary, new experimental data for the $^{27}\text{Al}(\text{p},\alpha)^{24}\text{Mg}$ rate appeared shortly after the CF88 compilation: a direct measurement by Timmermann et al. (1988) and a study of proton and alpha emission from ^{28}Si levels performed by Champagne et al. (1988). As a result, the rate is strongly reduced with respect to the Caughlan & Fowler (1988) one by up to 4 orders of magnitude in the region of interest. As a consequence, the calculated ^{26}Al yields were strongly reduced (José, Hernanz & Coc 1997).

The new rates available for $^{23}\text{Al}(\text{p},\gamma)^{24}\text{Si}$ (Schatz et al. 1997), $^{26}\text{Si}(\text{p},\gamma)^{27}\text{P}$ (Herndl et al. 1995), and $^{23}\text{Mg}(\text{p},\gamma)^{24}\text{Al}$ (Kubono, Kajino & Kato 1995 and Herndl et al. 1998) have no significant effect on the resulting ^{26}Al and ^{27}Al yields, as compared with the results obtained by José & Hernanz (1998) with $1.25 M_\odot$ ONe white dwarfs, using earlier prescriptions for these rates (i.e., van Wormer et al. 1994, Wiescher et al. 1986, and also Wiescher et al. 1986, respectively).

5. Updated nuclear reaction rates, nuclear uncertainties and their effect on the synthesis of ^{22}Na and ^{26}Al

Constraints on the synthesis of ^{22}Na and ^{26}Al during nova outbursts have been derived by means of hydrodynamic calculations involving ONe white dwarfs of masses 1.15, 1.25 and

$1.35 M_{\odot}$, assuming lower, recommended and upper estimates for several key reactions of the NeNa-MgAl cycles. Table 3 lists the set of reaction rates adopted for the calculation of minimum to maximum ^{22}Na and ^{26}Al production, compared with the rates adopted in José & Hernanz (1998). The specific choice adopted for these three sets is directly determined by the range of nuclear uncertainties accompanying the rates. The corresponding yields obtained with the different nuclear reaction networks, which are summarized in Tables 4 to 6, allow to derive error bars to the synthesis of ^{22}Na and ^{26}Al . Since we have not included other sources of uncertainty (convection, modelization of the explosion, ...) these error bars are only of nuclear physics origin.

5.1. Characteristics of the explosion

The early stages of the outburst are mainly dominated by the CNO cycle (José & Hernanz 1998). Therefore, the update of the nuclear reaction rates of the NeNa-MgAl cycles has no influence on the characteristics of the accretion phase (for instance, the duration of the accretion phase, t_{acc} , or the mass of the accreted envelope, ΔM_{env}).

Differences in the time evolution appear when the temperature near the burning shell reaches $T_{bs} \sim 10^8$ K. At this stage, energy generation by nuclear reactions involves a relevant contribution from the NeNa-MgAl cycles, together with the hot and cold modes of the CNO cycle. The use of updated rates modifies several properties of the TNR, such as the time required for a temperature rise from 3×10^7 K up to 10^8 K (hereafter, t_{rise}). For instance, a value of $t_{rise} = 6.8 \times 10^6$ s was found for Model ONe5 (José & Hernanz 1998), whereas a shorter time, $t_{rise} = 4.3 \times 10^6$ s, has been obtained in Model ONe125B, computed with updated (recommended) rates.

The role played by the NeNa-MgAl cycles increases when the temperature at the

location of the burning shell exceeds 2×10^8 K. This has an important effect on the energy production as well as on the resulting peak temperature achieved during the TNR. For instance, peak values for the nuclear energy generation rate and temperature achieved in Model ONe5, $\epsilon_{nuc,max} = 2.1 \times 10^{16} \text{ erg g}^{-1} \text{ s}^{-1}$ and $T_{max} = 2.44 \times 10^8$ K, translate into $\epsilon_{nuc,max} = 3.0 \times 10^{16} \text{ erg g}^{-1} \text{ s}^{-1}$ and $T_{max} = 2.51 \times 10^8$ K, for Model ONe125B. A similar trend was also pointed out by Starrfield et al. (1998), using updated nuclear reaction rates with respect to Caughlan & Fowler’s (1988) ones.

Such differences have also some influence on the last phases of the evolution, when the envelope expands, cools down and eventually a fraction of the formerly accreted shells is ejected. In general, we find that models computed with updated rates lead to slightly larger ejected masses with larger mean kinetic energies (see Tables 4–6) as a result of the larger $\epsilon_{nuc,max}$ attained. It is worth noticing that these differences in the properties of the TNRs will be reflected in the accompanying nucleosynthesis. The reason is twofold: first, the update of the reaction rates implies a certain modification of the nuclear paths. And second, the differences found in the time evolution as well as on the peak temperatures achieved modify the role played by (p, γ) and (p, α) reactions.

5.2. Nucleosynthesis in the NeNa-MgAl cycles

Due to the higher peak temperatures achieved in the models computed with the updated network (recommended rates. See Tables 4–6), ^{20}Ne is slightly reduced in the ejecta with respect to previous results obtained with older rates (José & Hernanz 1998). Differences are rather small (i.e., a mean mass fraction of 0.17 by mass, instead of 0.18, results for the 1.15 & 1.25 $M_{\odot}$ models), but they turn out to be important to determine the synthesis of less abundant species within the NeNa cycle. For instance, the abundance of ^{21}Ne increases by a factor between $\sim 3 - 5$ (the higher temperatures as well as the lower rate

adopted for $^{21}\text{Na}(\text{p},\gamma)^{22}\text{Mg}$ favor the chain $^{20}\text{Ne}(\text{p},\gamma)^{21}\text{Na}(\beta^+)^{21}\text{Ne}$, however no significant change is found for ^{22}Ne (which is essentially reduced by (p,γ) reactions from its initial amount). Worth noticing is also the net increase in the final amount of ^{22}Na in the ejecta: a factor of $\sim 4 - 5$ for the 1.15 & 1.25 $M_\odot$ models (resulting from the lower rates adopted for $^{21,22}\text{Na}(\text{p},\gamma)^{22,23}\text{Mg}$). This may have important consequences for gamma-ray astronomy, since it translates into a change by a factor of ~ 2 in the maximum expected distance at which the 1275 keV ^{22}Na line emitted by an exploding ONe novae would be eventually detected. Other isotopes, such as ^{23}Na and ^{25}Mg , are also overproduced (except for the 1.35 $M_\odot$ model), but less efficiently. As a result, the isotopic ratios $^{23}\text{Na}/^{22}\text{Na}$ decrease by nearly a factor of $\sim 2 - 3$. The abundances of the other magnesium isotopes, $^{24,26}\text{Mg}$, are slightly enhanced (in some cases even by a factor of $\sim 2 - 4$). Another interesting feature is that ^{26}Al remains essentially unaffected by the update of the network when the recommended rates are used. Therefore, the conclusions relative to the small contribution of classical nova outbursts to the synthesis of the Galactic ^{26}Al (see José, Hernanz & Coc 1997) still hold. Since the amount of ^{27}Al (and that of ^{28}Si) in the ejecta remains essentially unchanged, no variation on the isotopic ratio $^{26}\text{Al}/^{27}\text{Al}$ is found.

Another relevant outcome from the nucleosynthetic viewpoint is the dispersion in the mean ejected abundances obtained when the uncertainties associated with the nuclear reaction rates are taken into account. In the following, we will analyse the impact of such uncertainties in the resulting mean abundances in the ejecta. For that purpose, we will compare the yields obtained when the two extreme networks listed in Table 3 (i.e., A and C, leading to maximum and minimum ^{22}Na - ^{26}Al production, respectively) are adopted. We define the *dispersion factor*, F , as the ratio between the mean ejected abundances of a given nuclear species obtained when network A and C are adopted. All models computed (i.e., involving 1.15, 1.25 or 1.35 $M_\odot$ white dwarfs) show a very small dispersion in the ^{20}Ne yields, which results essentially from the different peak temperatures attained at the

envelope rather than from uncertainties in the nuclear reactions (such as $^{20}\text{Ne}(p,\gamma)^{21}\text{Na}$). No relevant dispersion is found either for ^{22}Ne , ^{23}Na , ^{27}Al , and ^{28}Si . On the contrary, the ^{21}Ne yields show a rather wide dispersion: a factor $F \sim 25$ in the $1.35 M_{\odot}$ models. Concerning ^{22}Na , differences between $F \sim 2 - 4$ are found. Similar degrees of dispersion are also obtained for the magnesium isotopes, with F ranging from 3 to 5 in the case of ^{25}Mg and from 2 to 3 for ^{24}Mg . Dispersion factors between 0.3 and 0.7 result for ^{26}Mg . The variation in the ejected amounts of ^{26}Al is particularly worth noticing, with a dispersion factor ranging from $F \sim 4$ to 7 (not considering the possible uncertainty associated with the 188 keV resonance in $^{26}\text{Al}^g(p,\gamma)^{27}\text{Si}$). Also interesting to notice is the fact that the $^{26}\text{Al}/^{27}\text{Al}$ ratio remains in the range 0.5–0.1 (see Tables 4–6).

In view of the abovementioned analysis, the nuclear uncertainties accompanying the reaction rates within the NeNa-MgAl cycles introduce a relatively wide dispersion in the yields resulting from classical nova outbursts. Therefore, predictions of ^{22}Na and ^{26}Al yields would benefit from new nuclear physics experiments aimed at reducing the uncertainties associated with some key reactions of the NeNa-MgAl cycles, in particular $^{21}\text{Na}(p, \gamma)^{22}\text{Mg}$ and $^{25}\text{Al}(p,\gamma)^{26}\text{Si}$ whose rates are uncertain by several orders of magnitude but also $^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$ and $^{22}\text{Na}(p, \gamma)^{23}\text{Mg}$. In addition, a verification of the yet unpublished values corresponding to the 0.188 MeV resonance of $^{26}\text{Al}^g(p,\gamma)^{27}\text{Si}$, as measured by Vogelaar (1989), is also recommended, due to its crucial role on the synthesis of ^{26}Al . These experiments should be focused on the measurement of a few key parameters (level energies, spectroscopic factors, ...). Many of these measurements involve short lived radioactive species and should benefit from the current development of radioactive ion beam facilities.

On the contrary, our calculations have shown that the remaining uncertainties on $^{22}\text{Mg}(p,\gamma)^{23}\text{Al}$, $^{23}\text{Al}(p,\gamma)^{24}\text{Si}$, $^{23}\text{Mg}(p,\gamma)^{24}\text{Al}$, $^{24}\text{Al}(p,\gamma)^{25}\text{Si}$, $^{26m}\text{Al}(p,\gamma)^{27}\text{Si}$, $^{26}\text{Si}(p,\gamma)^{27}\text{P}$, $^{26}\text{Mg}(p,\gamma)^{27}\text{Al}$, $^{27}\text{P}(p,\gamma)^{28}\text{S}$ or $^{27}\text{Si}(p,\gamma)^{28}\text{P}$ have no effect on ^{22}Na and ^{26}Al production.

6. Conclusions

We have computed a series of hydrodynamic models of nova outbursts, from the onset of accretion up to the ejection stage, for a range of ONe white dwarfs with masses between 1.15 and 1.35 $M_{\odot}$, with the aim of analyzing in detail the main nuclear paths leading to the synthesis of ^{22}Na and ^{26}Al . The role played by several key reactions within the NeNa-MgAl cycles has been tested by a series of models, which have been compared with previous results obtained by José & Hernanz (1998) with different prescriptions for the reaction rates. Limits on the production of both ^{22}Na and ^{26}Al have been derived from a series of computations assuming upper, recommended or lower estimates to the reaction rates. The most relevant conclusions extracted from this work can be summarized as follows:

1. The update of the nuclear reaction network results in a net increase in the final amount of ^{22}Na ejected into the interstellar medium during classical nova outbursts. This translates into an increase (by a factor of ~ 2) of the expected maximum distance at which an exploding ONe nova would be eventually detected through its emission at 1275 keV (^{22}Na decay γ -ray line).
2. The final amount of ^{26}Al remains essentially unaffected by the update of the network, confirming that classical novae scarcely contribute to the Galactic ^{26}Al , as pointed out earlier by José, Hernanz & Coc (1997).
3. Large nuclear uncertainties affect some key reactions of the NeNa-MgAl cycles, with a significant effect on the production of both ^{22}Na and ^{26}Al . When a combination of reaction rates, leading to maximum or minimum ^{22}Na - ^{26}Al synthesis, is adopted, a large dispersion in the final abundances is found. We stress that the derived ranges for ^{22}Na - ^{26}Al production can be interpreted as error bars on sodium & aluminum production posed by nuclear physics uncertainties.

4. In order to reduce the impact of the nuclear uncertainties in the production of ^{22}Na and ^{26}Al , we point out some nuclear reactions that deserve new experiments, in particular, $^{21}\text{Na}(\text{p}, \gamma)^{22}\text{Mg}$, $^{22}\text{Na}(\text{p}, \gamma)^{23}\text{Mg}$ and $^{25}\text{Al}(\text{p}, \gamma)^{26}\text{Si}$. A confirmation of the values corresponding to the 0.188 MeV resonance of $^{26}\text{Al}^g(\text{p}, \gamma)^{27}\text{Si}$, as measured by Vogelaar (1989), would also be of great importance.

A. Appendix: Nuclear reactions

We discuss here the reaction rates corresponding to proton capture reactions on short-lived nuclei. Other reactions within the NeNa and MgAl cycles are analyzed in the text (see Sections 3.2 and 4.2) and more extensively in the forthcoming *Compilation of Charged-Particle Induced Thermonuclear Reaction Rates* (Angulo et al. 1998). First, we introduce some standard nuclear physics quantities and notations to be used hereafter. We also briefly summarize several indirect methods that are generally used to extract or estimate unavailable direct data, for the calculation of thermonuclear rates.

For an $A(x, y)B$ reaction, resonance strengths (at $E = E_r$) are given by $\omega\gamma$, where $\omega = (2J_r + 1)((2J_x + 1)(2J_A + 1))^{-1}$ is the statistical spin factor and $\gamma = \Gamma_x\Gamma_y/\Gamma$ is the width ratio. The total width ($\Gamma(E)$) corresponds to the sum of partial widths ($\Gamma_I(E)$) : $\Gamma = \sum_{I=x,y,\dots} \Gamma_I$. The spectroscopic factor, C^2S , relates single particle widths ($\Gamma_x^{s.p.}$) to actual ones: $\Gamma_x = C^2S \Gamma_x^{s.p.}$. These single particle widths are calculated by solving the Schrödinger equation for a particle in a realistic nuclear potential. An approximation for the upper limit of single particle widths is given by the *Wigner limit*: $\Gamma_x^W = 3\hbar^2(\mu s^2)^{-1}P_{\ell_x}(E = E_r, s)$, where s is the channel radius, μ the reduced mass, $P_{\ell_x}(E, s)$ the penetration factor and ℓ_x the orbital angular momentum transferred by particle x . We introduce the reduced width, θ_x^2 , defined by $\Gamma_x = \theta_x^2\Gamma_x^W$, as it is sometimes used to derive estimates of reaction rates (see Iliadis 1997 for a discussion of the relationship between θ^2 and C^2S).

Experimental spectroscopic factors can be extracted from transfer reactions (e.g., the proton transfer reaction $X(^3\text{He}, d)Y$ will lead to one proton spectroscopic factors used to calculate proton widths entering into the calculation of the $X(p, \gamma)Y$ rate). Spectroscopic factors for conjugate reactions (obtained by the $p \rightleftharpoons n$ exchange) are assumed to be approximately equal. For instance, neutron spectroscopic factors for ^{22}Ne

can be extracted from the one neutron $^{21}\text{Ne}(\text{d},\text{p})^{22}\text{Ne}$ transfer reaction. According to the above approximation, these *neutron* spectroscopic factors can be used to determine *proton* widths for the calculation of the $^{21}\text{Na}(\text{p},\gamma)^{22}\text{Mg}$ rate for instance. Radiative widths (Γ_γ) corresponding to transitions of the magnetic ($M\lambda$) or electric type ($E\lambda$) of order λ are often expressed in Weiskopf's units (W.u.) to remove the effect of their $E_\gamma^{2\lambda+1}$ dependence and strong variation with λ (e.g. Firestone et al. 1996). Compilations of Γ_γ , expressed in these units, are available (Endt 1979) and may be used to estimate unknown radiative widths. They can also be obtained from the conjugate nuclei after correction for the $E_\gamma^{2\lambda+1}$ dependence. If particle emission is energetically forbidden in the conjugate level, the radiative width equals the total width. If the lifetime of this level is known, the radiative width is easily obtained.

A.1. The $^{21}\text{Na}(\text{p},\gamma)^{22}\text{Mg}$ reaction

Estimates of the $^{21}\text{Na}(\text{p},\gamma)^{22}\text{Mg}$ reaction rate (Caughlan & Fowler 1988) have been provided by Wiescher et al. (1986) and Wiescher & Langanke (1986), considering the first three levels (Endt 1990) above the proton threshold. With the exception of the first level, the proton widths are much larger than the gamma widths, so that $\omega_p \approx \omega_\gamma$. Only the first two levels ($E_x=5.714$ and 5.837 MeV) contribute to the rate in the temperature domain considered. The $E_x = 5.837$ MeV level is assumed to be the conjugate of the $E_x = 5.910$ one in ^{22}Ne (Endt 1990). If this assignment is correct, the corresponding gamma width can be reliably deduced (Wiescher & Langanke 1986). On the contrary, the strength of the first $E_x=5.714$ MeV, $J^\pi=2^+$ level suffers from a significant uncertainty. For this level, the total width is known experimentally to be $\Gamma=16.5\pm4.4$ meV. To calculate the corresponding resonance strength, Wiescher & Langanke (1986) and Wiescher et al. (1986) estimated the proton width, Γ_p , and deduced the radiative width, Γ_γ , from the relation

$\Gamma = \Gamma_p + \Gamma_\gamma$. To estimate the proton width, two hypotheses were considered: Wiescher et al. (1986) assumed $\ell_p = 0$ and $\theta_p^2 = 0.01$, but later Wiescher & Langanke (1986) took $\ell_p = 2$ and $\theta_p^2 = 0.5$. Nevertheless, because of the $\Gamma_p + \Gamma_\gamma = 16.5 \pm 4.4$ meV constraint, the corresponding width ratios, γ , are not much different: 3.4 or 3.8 meV, very close to the maximum value ($\Gamma/4$), obtained when $\Gamma_\gamma = \Gamma_p = \Gamma/2$. There are two reasons to think that, on the contrary, $\Gamma_p \ll \Gamma \approx \Gamma_\gamma$ and that the corresponding strength is much smaller than the estimates provided by Wiescher et al. First, this level has a known counterpart in ^{22}Ne at $E_x = 6.120$ MeV (Endt 1998) with a measured, purely radiative, total width $\Gamma = 29 \pm 9$ meV, that enables to calculate the radiative width of the $E_x = 5.714$ MeV ^{22}Mg level to be $\Gamma_\gamma \simeq 23$ meV. Since it is found to be greater than the measured total width (16.5 ± 4.4 meV), it is likely that $\Gamma_p \ll \Gamma$ for this level. The second indication comes from experimental data on the $^{21}\text{Ne}(d,p)^{22}\text{Ne}$ reaction, which provides information on neutron capture on ^{21}Ne (i.e., the mirror counterpart of proton capture on ^{21}Na). Neutron spectroscopic factors were obtained by Neogy et al. (1972) for many ^{22}Ne levels, from $E_x = 0$ to 9.07 MeV, but not for the $E_x = 6.120$ MeV one due to its very low population and flat angular distribution. Indeed, the direct comparison of peak heights in the experimental spectrum (fig. 1, Neogy et al. 1972) shows that, within levels of same spin and transferred angular momentum, the cross section for the production of the $E_x = 6.120$ MeV level is very small. In addition, the corresponding angular distribution (fig. 5, Neogy et al. 1972) is very flat, indicating no evidence for a direct component and hence a very small neutron spectroscopic factor. This conclusion can be extended to the proton spectroscopic factor in the ^{22}Na mirror level. The 6.551 MeV ^{22}Na level, is assumed (Endt 1990, Endt 1998) to belong to the same isospin triplet as the 6.120 MeV ^{22}Ne level. Its proton spectroscopic factor has been measured by Garrett et al. (1971) to be ≈ 0.1 . This value is not consistent with the Neogy et al. (1972) data and would result, for the 5.714 MeV ^{22}Mg level, in a proton width in excess of the measured, 16.5 meV, total width. Hence one may question the isospin triplet assignment

made by Endt (1990) and Endt (1998). Moreover, the 5.714 MeV ^{22}Mg level is withdrawn from the triplet in the last paper by Endt (1998). Accordingly, we consider the Wiescher et al. estimate for this resonance strength as an upper limit and adopt $\omega\gamma = 2.5, 0.25, 0.0$ meV for upper ($\Gamma_\gamma = \Gamma_p = \Gamma/2$), recommended value (with the usual 0.1 reduction factor) and lower limit. We provide below the updated formula for the rate including the uncertainty factors and the direct capture contribution from Caughlan & Fowler (1988):

$$\begin{aligned}
 & 1.41 \times 10^5 T_9^{-3/2} \exp\left(-20.739/T_9^{1/3} - (T_9/.366)^2\right) \times \\
 & (1. + .02 T_9^{1/3} + 4.741 T_9^{2/3} + 667 T_9 + 16.38 T_9^{4/3} + 5.858 T_9^{5/3}) + \quad DC \text{ (CF88)} \\
 & (0 \rightarrow 10.) \times 40.8. T_9^{-3/2} \exp(-2.52/T_9) + \quad (E_x = 5.714) \\
 & 1857. T_9^{-3/2} \exp(-3.95/T_9) + \quad (E_x = 5.837) \\
 & (0.1 \rightarrow 10.) \times 408. T_9^{-3/2} \exp(-5.49/T_9) \quad (E_x = 5.965)
 \end{aligned}$$

The third term corresponds to the 5.837 MeV level, assuming that the analog assignment made by Endt (1990) is correct. This is however not granted as it is not present anymore in Endt (1998). The last term corresponds to the $E_x=5.965$ MeV, $J^\pi = 0^+$ level. Since its widths are unknown (save that $\Gamma_\gamma \ll \Gamma \approx \Gamma_p$), Wiescher et al. (1986) assumed a typical value for Γ_γ based on the statistics of Endt (1979). We introduce a factor of ten uncertainty to account for the dispersion of the gamma strengths within this statistics, but it has a little influence on the rate. The corresponding rates, relative to Caughlan & Fowler (1988) (i.e., Wiescher et al. 1986) are shown in Figure 4: the maximum effect occurs between $T_8 = 0.5$ and 3, well within the nova temperature range. Experimental data of the $E_x=5.714$ MeV, $J^\pi=2^+$ level is clearly needed to improve the reliability of this rate.

A.2. The $^{22}\text{Mg}(\text{p},\gamma)^{23}\text{Al}$ and $^{23}\text{Al}(\text{p},\gamma)^{24}\text{Si}$ reactions

As ^{22}Mg plays a crucial role on the synthesis of ^{22}Na through its beta decay, its destruction by $^{22}\text{Mg}(\text{p},\gamma)^{23}\text{Al}$ has to be considered. The contribution of the direct capture to the ground state for the $^{22}\text{Mg}(\text{p},\gamma)^{23}\text{Al}$ rate has been calculated by Wiescher et al. (1986). Later on, Wiescher et al. (1988) measured the location of the first resonance and also calculated its strength (shell model and mirror level). But due to the very low Q-value (0.125 MeV), rapid photodisintegration of ^{23}Al prevents ^{22}Mg destruction. Another destruction channel has also been proposed: it takes place via two proton captures on ^{22}Mg (i.e., proton capture on the small population of ^{23}Al through $^{23}\text{Al}(\text{p},\gamma)^{24}\text{Si}$. Görres, Wiescher & Thielemann 1995) and a new rate was proposed by Herndl et al. (1995), based on shell model calculations. The location of the first two ^{24}Si levels has been recently determined experimentally (Schatz et al. 1997): it leads to a higher rate than in Herndl et al. (1995) due to the lower location of the first resonance ($E_r^{cm} = 0.141 \pm 0.031$ MeV, instead of 320 keV). In our nova models, we have adopted the new limits for the $^{23}\text{Al}(\text{p},\gamma)^{24}\text{Si}$ rate given by Schatz et al. (1997).

A.3. The $^{23}\text{Mg}(\text{p},\gamma)^{24}\text{Al}$ and $^{24}\text{Al}(\text{p},\gamma)^{25}\text{Si}$ reactions

An estimate to the $^{23}\text{Mg}(\text{p},\gamma)^{24}\text{Al}$ rate has been provided by Wiescher et al. (1986) based on direct capture and three resonances. Since then, Kubono, Kajino & Kato (1995), Endt (1998) and Herndl et al. (1998) have discussed analog level assignments, in particular with the help of shell model calculations and isobaric multiplet mass equation (Herndl et al. 1998). For nova nucleosynthesis, only the two first resonances have to be considered. They correspond to the $E_x = 2.349$ and 2.534 MeV levels and are assumed to be the analogs of the 2.514 MeV, 3^+ and 2.563 MeV, $4^+(2^+)$ ones in ^{24}Na (Endt 1998, Herndl et al. 1998). In addition to the estimate provided by Wiescher et al. (1986) the strength of the second

resonance has been calculated (shell model) by Herndl et al. 1998. However, combining statistics of reduced radiative widths (Endt 1979) together with the spectroscopic factor reported in Endt & Van Der Leun (1978), show that its contribution is, in any case, negligible with respect to the first one in the domain of temperature considered here. The strength of the first resonance is deduced from the analog level ($\Gamma_\gamma \ll \Gamma_p$) and hence suffers only small uncertainty like the direct capture contribution (see Wiescher et al. 1986). In the domain of nova nucleosynthesis, the main uncertainty comes from the determination of the energy of the first level: taking $E_r^{cm} = 0.458$ (Kubono et al. 1995), 0.478 (Endt 1998; Herndl et al. 1998) or 0.51 MeV (Wiescher et al. 1986) results in a variation of the rate of less than a factor of ~ 10 around $T_8 = 2.5$. To check the importance of this reaction, we adopted the highest rate (Kubono et al. 1995) in our test calculations.

For $^{24}\text{Al}(\text{p},\gamma)^{25}\text{Si}$, we use the rate proposed by Herndl et al. (1995), based on shell model calculations. It remains very uncertain, due to the limited experimental spectroscopic information on ^{25}Si . It should have, however, very little effect on Ne–Na leaks, since either $^{24}\text{Al}(\beta^+)^{24}\text{Mg}(\text{p},\gamma)^{25}\text{Al}$ or $^{24}\text{Al}(\text{p},\gamma)^{25}\text{Si}(\beta^+)^{25}\text{Al}$ lead to ^{25}Al in the MgAl cycle.

A.4. The $^{25}\text{Al}(\text{p},\gamma)^{26}\text{Si}$ reaction

$^{25}\text{Al}(\text{p},\gamma)^{26}\text{Si}$ is important since it leads to the formation of the short-lived isomer ($^{26}\text{Al}^m$) instead of the long-lived ground state ($^{26}\text{Al}^g$). Its rate suffers from large uncertainties in the domain of nova nucleosynthesis due mainly to the unknown location of the analog of the ^{26}Mg $E_x = 6.13$ MeV, 3^+ level. A level shift similar to the one of its immediate neighbor ($E_x = 6.26$ MeV; 0^+ in ^{26}Mg) would bring the 3^+ level within the Gamow peak with a dramatic influence on the rate. The maximum effect on ^{26}Al production is found when $E_r^{cm}(3^+) = 0.2 \pm 0.1$ MeV (Coc et al. 1995). New Coulomb displacement energy calculations (Iliadis et al. 1996) give $E_r^{cm}(3^+) = 0.45 \pm 0.1$ MeV. However, discrepancies

between calculated and experimentally known ^{26}Si level energies can exceed ± 0.1 MeV so that the case $E_r^{cm}(3^+) = 0.2$ cannot be ruled out. (See, e.g., the $^{23}\text{Al}(p,\gamma)^{24}\text{Si}$ case above where the location of the first resonance is found $\simeq 180$ keV below the calculated one.) Accordingly, for this rate, we used the lower limit, recommended value and higher limit provided by Coc et al. (Case A, B and C, respectively).

A.5. The $^{26}\text{Si}(p,\gamma)^{27}\text{P}$ reaction

The $^{26}\text{Si}(p,\gamma)^{27}\text{P}$ reaction rate has been discussed by Wiescher et al. (1986), and more recently by Herndl et al. (1995). Only two excited states are known in ^{27}P and only one ($E_x = 1.660$ MeV, $J^\pi = 5/2^+$) is low enough to be of astrophysical interest. From the comparison with the spectrum of the mirror nucleus, another level, $J^\pi = 3/2^+$, located at $E_x = 0.985$ MeV in ^{27}Mg , is expected below the $5/2^+$ one. Wiescher et al. (1986) assumed a small shift, while shell model calculations (Herndl et al. 1995) led to a much larger shift, which resulted in a dramatic increase of the rate by almost four orders of magnitude. The calculated strength (Herndl et al. 1995) is in good agreement with the Γ_γ value (0.8 meV) inferred from the lifetime of the $E_x = 0.985$, ^{27}Mg , conjugate level and with the Γ_p obtained from the neutron spectroscopic factor in the conjugate ^{27}Mg level reported in Endt & Van Der Leun (1978). In consequence, the uncertainty results from the unknown position of the resonance. However, it does not affect strongly the rate (a factor of ten at most) for $1.5 < T_8 < 4$ down to $E_r^{cm} \lesssim 0.2$ MeV due to the compensating effect of the evolution of the penetrability $P_{l_p}(E_r, s)$ and $\exp(-E_r/kT)$ factors with E_r small variations. The strength of the $5/2^+$ resonance is known from the Wiescher et al. (1986) estimates, from ^{27}Mg spectroscopy and from the Herndl et al. (1995) calculations. However, its location ($E_r = 0.76$ MeV) makes its contribution negligible compared to the direct capture one at nova temperatures. The latter provides the lower limit for the rate, reflecting the possibility that

the $J^\pi = 3/2^+$ resonance energy is lower than $E_r^{cm} \simeq 0.2$ MeV, as considered by Wiescher et al. (1986), while the upper limit is given by the Herndl et al. (1995) rate. One should note that this effect could be enhanced if the uncertain $^{25}\text{Al}(p,\gamma)^{26}\text{Si}$ rate is close to its maximum value (see above).

A.6. The $^{27}\text{P}(p,\gamma)^{28}\text{S}$, and $^{27}\text{Si}(p,\gamma)^{28}\text{P}$ reactions

Only the ground state of ^{28}S is known but from the level scheme of the conjugate nucleus, ^{28}Mg , one can infer that no resonance is expected for $E_r^{cm} \lesssim 1.4$ MeV. Hence, at the temperatures considered in the text, the $^{27}\text{P}(p,\gamma)^{28}\text{S}$ reaction proceeds only through direct capture. The corresponding rate has been obtained by Herndl et al. (1995) from shell model calculations. It is about a factor of ten higher than the Caughlan & Fowler (1988) one and can be considered as reliable enough (i.e. within an order of magnitude), due to the expected absence of resonances.

The $^{27}\text{Si}(p,\gamma)^{28}\text{P}$ reaction rate given in Caughlan & Fowler (1998) comes from the analysis of Wiescher et al. (1986). They considered five resonances corresponding to the ^{28}P levels between $E_x = 2.143$ and 2.628 MeV, out of which, for three levels, the spins are uncertain. From the assumed conjugate levels in ^{28}Al , they extracted radiative widths and proton widths using spectroscopic factors obtained by neutron transfer on ^{27}Al . More recently, the good agreement between calculated (Endt & Booten 1993) and experimental (Endt 1990) level schemes, spectroscopic factors and radiative widths, for $A = 28$ nuclei, gives confidence to the ^{28}Al and ^{28}P conjugate level assignment (see also Iliadis et al. 1998). The first level above proton threshold, $E_x = 2.104$ MeV, $J^\pi = 2^+$, was not considered by Wiescher et al. (1986), but its estimated proton width is however too small ($\Gamma_p \lesssim \Gamma_p^W \simeq 10^{-20}$ eV) to contribute to the rate for $T_8 \gtrsim 0.3$. The uncertainty remains limited to transposition of ^{28}Al data to ^{28}P .

The authors are grateful to Jean-Pierre Thibaud for helpful and stimulating discussions. We also thank Carmen Angulo for allowing us to use information from the NACRE compilation prior to publication, and the referee, Sumner Starrfield, for a critical reading of the manuscript. This research has been partially supported by the CICYT-P.N.I.E. (ESP98-1348), by the DGICYT (PB97-0983-C03-02; PB97-0983-C03-03), by the CIRIT (GRQ94-8001), by the AIHF1997-0087, by the CHRXCT930339 and by the PICS 319.

REFERENCES

- Angulo, C. et al. 1998 (The NACRE collaboration), Nucl. Phys. A, in press
- Arnett, W.D. & Truran, J.W. 1969, ApJ, 157, 339
- Caughlan, G.R. & Fowler, W.A. 1988, At. Data Nucl. Data Tables, 40, 283
- Champagne, A.E. et al. 1988, Nucl. Phys. , A487, 433
- Champagne, A.E., Magnus, P.V., Smith, M.S. & Howard, A.J. 1990, Nucl. Phys. , A512, 317
- Clayton, D.D. & Hoyle, F. 1974, ApJ, 187, L101
- Coc, A., Mochkovitch, R., Oberto, Y., Thibaud J.-P. & Vangioni-Flam, E. 1995, A&A 299, 479
- Coc, A. & Porquet, M.G. 1998, in Proc. Tours Symposium on Nuclear Physics III, ed. M. Arnould, M. Lewitowicz, Yu. Ts. Oganessian, M. Ohta, H. Utsonomiya & T. Wada (New York:AIP), p. 457
- DeWitt, H.E., Graboske, H.C. & Cooper, M.S. 1973, ApJ, 181, 439
- Diehl, R. et al. 1995, A&A, 298, 445
- Diehl, R. et al. 1997, in Proc. Fourth Compton Symposium, ed. C.D. Dermer, M.S. Strickman & J.D. Kurfess (New York:AIP), p. 1114
- Domínguez, I., Tornambé, A. & Isern, J. 1993, ApJ, 419, 268
- Endt, P.M. 1979, At. Data Nucl. Data Tables, 23, 3
- Endt, P.M. 1990, Nucl. Phys. , A521, 1

- Endt, P.M. & Booten, J.G.L. 1993, Nucl. Phys. , A555, 499
- Endt, P.M. & Rolfs, C. 1987, Nucl. Phys. A467, 261
- Endt, P.M. & Van Der Leun, C. 1978, Nucl. Phys. , A301, 1
- Endt, P.M. 1998, Nucl. Phys. , A633, 1
- Firestone, R.B., Shirley, V.S., Baglin, C.M., Chu, S.Y.F. & Zipkin, J. 1996, Table of Isotopes, John Wiley & Sons ed.
- Garrett, J.D., Middleton, R., Fortune, H.T. 1971, Phys. Rev., C4, 165
- Gehrz, R.D., Truran, J.W., Williams, R.E. & Starrfield, S. 1998, Publ. Astron. Soc. Pacific, 110, 3
- Gómez–Gomar, J., Hernanz, M., José, J. & Isern, J. 1998, MNRAS, 296, 913
- Görres, J. et al. 1983, Nucl. Phys. , A408, 372
- Görres, J., Wiescher, M. & Rolfs, C. 1989, ApJ , 343, 365
- Görres, J., Wiescher, M. & Thielemann, F.-K. 1995, Phys. Rev., C51, 392
- Graboske, H.C., DeWitt, H.E., Grossman, A.S. & Cooper, M.S. 1973, ApJ, 181, 457
- Hernanz, M., Gómez–Gomar, J., José, J. & Isern, J. 1997, in 2nd INTEGRAL Workshop ‘The Transparent Universe’, ed. C. Winkler, T.J.L. Courvoisier & Ph. Durouchoux (Noordwijk:ESA SP-382), 47
- Herndl, H. et al. 1995, Phys. Rev., C52, 1078
- Herndl, H., Fantini, M., Iliadis, C., Endt, P.M., & Oberhummer, H. 1998, Phys. Rev. C58, 1798

- Hillebrandt, W. & Thielemann, F.-K. 1982, ApJ, 225, 617
- Iliadis, C. 1997, Nucl. Phys. , A618, 166
- Iliadis, C. et al., 1990, Nucl. Phys. , A512, 509
- Iliadis, C., Buchmann, L., Endt, P.M., Herndl, H. & Wiescher, M. 1996, Phys. Rev., C53, 475
- Iliadis, C., Endt, P.M., Prantzos, N. & Thompson, W.J. 1998, ApJ, in press
- Iyudin, A.F. et al. 1995, A&A, 300, 422
- José, J. 1996, PhD Thesis, University of Barcelona
- José, J. & Hernanz, M. 1997, Nucl. Phys., A621, 491
- José, J. & Hernanz, M. 1998, ApJ , 494, 680
- José, J., Hernanz, M. & Coc, A. 1997, ApJ , 479, L55
- Knödlseeder, J. 1997, PhD Thesis, Paul Sabatier University (Toulouse)
- Kovetz, A. & Prialnik, D. 1997, ApJ, 477, 356
- Kubono, S., Kajino, T., & Kato, S., 1995, Nucl. Phys. , A588, 521
- Leising, M.D. 1993, A&AS , 97, 299
- Leising, M.D., Share, G.H., Chupp, E.L. & Kanbach, G. 1988, ApJ, 328, 755
- Leventhal, M., MacCallum, C. & Watts, A. 1977, ApJ, 216, 491
- Livio, M., Truran, J.W. 1994, ApJ, 425, 797
- Mahoney, W.A., Ling, J.C., Jacobson, A.S. & Lingenfelter, R.E. 1982, ApJ, 262, 742

- Mahoney, W.A., Ling, J.C., Wheaton, W.A. & Jacobson, A.S. 1984, ApJ, 286, 578
- Neogy, P., Middleton, R. & Scholtz, W. 1972, Phys. Rev., C6, 885
- Nofar, I., Shaviv, G. & Starrfield, S. 1991, ApJ, 369, 440
- Politano, M., Starrfield, S., Truran, J.W., Weiss, A. & Sparks W.M. 1995, ApJ, 448, 807
- Prantzos, N. & Diehl, R., 1996, Phys. Rep., 267, 1
- Prialnik, D., Shara, M.M. & Shaviv, G. 1979, A&A, 72, 192
- Ritossa, C., García-Berro, E. & Iben, I. 1996, ApJ, 460, 489
- Rolfs, C., Rodney, W.S., Shapiro, M.H., & Winkler H., 1975 Nucl. Phys. , A241, 460
- Schatz, H. et al. 1997, Phys. Rev. Lett., 79, 3845
- Schmidt, S. et al. 1995, Nucl. Phys. , A591, 227
- Seuthe, S. et al. 1990, Nucl. Phys. , A514, 471
- Share, G.H., Kinzer, R.L., Kurfess, J.D., Forrest, J.D., Chupp, F.L. & Rieger, E. 1985, ApJ, 292, L61
- Starrfield, S., Truran, J.W., Sparks, W.M. & Kutter, G.S. 1972, ApJ, 176, 169
- Starrfield, S., Truran, J.W., Wiescher, M.C. & Sparks, W.M. 1997, Nucl. Phys., A621, 495
- Starrfield, S., Truran, J.W., Wiescher, M.C. & Sparks, W.M. 1998, MNRAS, 296, 502
- Stegmüller, F. et al. 1996, Nucl. Phys. , A601, 168
- Timmermann, R., Becker, H.W., Rolfs, C., Schröder, U. & Trautvetter, H.P. 1988, Nucl. Phys., A477, 105

- Van Wormer, L., Görres, J., Iliadis, C., Wiescher, M. & Thielemann, F.-K. 1994, ApJ , 432
326
- Vogelaar R.B. 1989, PhD Thesis, California Institute of Technology
- Vogelaar R.B. et al. 1996, Phys. Rev, C53, 1945
- Wallace, R.K. & Woosley, S.E. 1980, ApJS, 45, 247
- Ward, R.A. & Fowler, W.A. 1980, ApJ , 238, 266
- Weiss, A. & Truran, J.W. 1990, A&A, 238, 178
- Wiescher, M. & Langanke, K. 1986, Z. Phys., 325, 309
- Wiescher, M., Görres, J., Thielemann, F.-K. & Ritter, H. 1986, A&A , 160, 56
- Wiescher, M., Görres, J., Sherrill, B., Mohar, M., Winfield, J.S. & Brown, B.A. 1988, Nucl.
Phys. , A484, 90
- Wood, P.R. 1974, ApJ, 190, 609

Table 1. Initial composition of the envelope (up to Si), assuming 50% mixing with the

ONe white dwarf core

Nuclei	Mass fraction
^1H	3.5E-1
^3He	1.5E-5
^4He	1.4E-1
^6Li	3.2E-10
^7Li	4.7E-9
^9Be	8.3E-11
^{10}B	5.3E-10
^{11}B	2.4E-9
^{12}C	6.1E-3
^{13}C	1.8E-5
^{14}N	5.5E-4
^{15}N	2.2E-6
^{16}O	2.6E-1
^{17}O	1.9E-6
^{18}O	1.1E-5
^{19}F	2.0E-7
^{20}Ne	1.6E-1
^{21}Ne	3.0E-3
^{22}Ne	2.2E-3
^{23}Na	3.2E-2
^{24}Mg	2.8E-2
^{25}Mg	7.9E-3
^{26}Mg	5.0E-3

Table 1—Continued

Nuclei	Mass fraction
^{27}Al	5.4E-3
^{28}Si	3.3E-4

Table 2. Test models

Nuclear reaction	Old rate	Test rate	$\frac{X(^{22}\text{Na})_{\text{test}}}{X(^{22}\text{Na})_{\text{old}}}$	$\frac{X(^{26}\text{Al})_{\text{test}}}{X(^{26}\text{Al})_{\text{old}}}$
1.15 $M_{\odot}$ ONe				
$^{21}\text{Na}(p, \gamma)^{22}\text{Mg}$	CF88	CF88/100	2.3	1.1
$^{25}\text{Al}(p, \gamma)^{26}\text{Si}$	Wie86	Coc95, case A	1	1.2
	Wie86	Coc95, case C	1	0.5
$^{23}\text{Na}(p, \gamma)^{24}\text{Mg}$	CF88	CF88+GWR89	1.2	1.3
$^{26}\text{Al}^g(p, \gamma)^{27}\text{Si}$	Vog89	Vog89	1	1.9
1/3 $\times$ res(0.188 MeV)				
$^{26}\text{Al}^m(p, \gamma)^{27}\text{Si}$	CF88	CF88 $\times$ 100	1	1
$^{26}\text{Si}(p, \gamma)^{27}\text{P}$	Wie86	Her95	1	1
$^{26}\text{Mg}(p, \gamma)^{27}\text{Al}$	Ili90	Ili90+Cha90	1	1
1.25 $M_{\odot}$ ONe				
$^{21}\text{Na}(p, \gamma)^{22}\text{Mg}$	CF88	CF88/100	3	1.1
$^{22}\text{Na}(p, \gamma)^{23}\text{Mg}$	CF88	Ste96	3	1.2
$^{23}\text{Mg}(p, \gamma)^{24}\text{Al}$	Wie86	KKK95	1.1	1.1
1.35 $M_{\odot}$ ONe				
$^{21}\text{Na}(p, \gamma)^{22}\text{Mg}$	CF88	CF88/100	1.2	1
$^{23}\text{Al}(p, \gamma)^{24}\text{Si}$	Wor94	Sch97	1	1

References to the reaction rates: Wie86 (Wiescher et al. 1986), CF88 (Caughlan & Fowler 1988), GWR89 (Görres, Wiescher & Thielemann 1989), Vog89 (Vogelaar 1989), Cha90 (Champagne et al. 1990), Ili90 (Iliadis et al. 1990), Wor94 (van Wormer et al. 1994), Coc95 (Coc et al. 1995), Her95 (Herndl et al. 1995), KKK95 (Kubono, Kajino & Kato 1995), Ste96 (Stegmüller et al. 1996), Sch97 (Schatz et al. 1997)

Table 3. Reaction rates adopted for the calculation of Maximum (A), Recommended (B), and Minimum (C) ^{22}Na - ^{26}Al production compared with those adopted in José & Hernanz (1998)

Reaction	JH98	A	B	C
$^{21}\text{Na}(p,\gamma)^{22}\text{Mg}$	CF88	This work Lower limit	This work Recommended	This work Upper limit
$^{21}\text{Ne}(p,\gamma)^{22}\text{Na}$	CF88 with f=0.1	CF88 f=0.01	CF88 f=0.001	CF88 f=0
$^{22}\text{Na}(p,\gamma)^{23}\text{Mg}$	CF88	Ste96 Lower limit	NACRE Recommended	Ste96 Upper limit
$^{25}\text{Al}(p,\gamma)^{26}\text{Si}$	Wie86	Coc95 Case A (Lower)	Coc95 Case B (Recommended)	Coc95 Case C (Upper)
$^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$	CF88	CF88+GWR89 with f=1	CF88+GWR89 f=0.1	CF88+GWR89 f=0
$^{23}\text{Mg}(p,\gamma)^{24}\text{Al}$	Wie86	KKK95	KKK95	KKK95
$^{26}\text{Mg}(p,\gamma)^{27}\text{Al}$	Ili90	Cha90+Ili90 with f=1	Cha90+Ili90 f=0.1	Cha90+Ili90 f=0
$^{26}\text{Al}^m(p,\gamma)^{27}\text{Si}$	CF88	CF88	CF88	CF88
$^{27}\text{Al}(p,\alpha)^{24}\text{Mg}$	Cha88	Cha88+Tim88 with f=1	Cha88+Tim88 f=0.1	Cha88+Tim88 f=0
$^{23}\text{Al}(p,\gamma)^{24}\text{Si}$	WW80	Sch97 Lower limit	Sch97 Recommended	Sch97 Upper limit
$^{26}\text{Si}(p,\gamma)^{27}\text{P}$	Wie86	Her95	Her95	Her95
$^{26}\text{Al}^g(p,\gamma)^{27}\text{Si}$	Vog89	Coc95 with f1=f2=0	Coc95 f1=f2=0.1	Coc95 f1=f2=1

Additional references to the reaction rates: WW80 (Wallace & Woosley 1980), Cha88 (Champagne et al. 1988), Tim88 (Timmermann et al. 1988), NACRE (Angulo et al. 1998). f, f1 and f2 represent uncertainty factors in the analytic rates.

Table 4. Main characteristics of the 1.15 $M_{\odot}$ ONe nova models compared with model ONe3 (José & Hernanz 1998)

	ONe3	ONe115A	ONe115B	ONe115C
Reaction Network	Old	A	B	C
ΔM_{env} ($10^{-5} M_{\odot}$)	3.2	3.2	3.2	3.2
T_{max} (10^8 K)	2.19	2.33	2.31	2.30
ΔM_{ejec} ($10^{-5} M_{\odot}$)	1.9	2.6	2.6	2.6
K (10^{45} erg)	1.2	1.5	1.5	1.5
^{20}Ne	1.8E-1	1.6E-1	1.7E-1	1.8E-1
^{21}Ne	3.0E-5	2.0E-4	8.7E-5	2.5E-5
^{22}Ne	1.7E-3	1.6E-3	1.5E-3	1.5E-3
^{22}Na	5.3E-5	5.3E-4	2.7E-4	1.2E-4
^{23}Na	7.5E-4	1.3E-3	1.4E-3	1.4E-3
^{24}Mg	1.0E-4	9.6E-4	4.1E-4	3.5E-4
^{25}Mg	2.9E-3	6.1E-3	3.9E-3	1.8E-3
^{26}Mg	3.4E-4	8.3E-4	6.4E-4	1.2E-3
^{26}Al	9.3E-4	1.2E-3	8.0E-4	3.3E-4
^{27}Al	4.5E-3	4.8E-3	3.5E-3	3.4E-3
^{28}Si	5.4E-2	6.8E-2	5.8E-2	5.6E-2

Table 5. Main characteristics of the 1.25 M_⊙ ONe nova models compared with model ONe5 (José & Hernanz 1998)

	ONe5	ONe125A	ONe125B	ONe125C
Reaction Network	Old	A	B	C
ΔM_{env} (10 ⁻⁵ M _⊙)	2.2	2.2	2.2	2.2
T_{max} (10 ⁸ K)	2.44	2.54	2.51	2.51
ΔM_{ejec} (10 ⁻⁵ M _⊙)	1.4	1.8	1.8	1.8
K (10 ⁴⁵ erg)	1.4	1.5	1.5	1.5
²⁰ Ne	1.8E-1	1.6E-1	1.7E-1	1.7E-1
²¹ Ne	3.5E-5	3.7E-4	1.4E-4	3.0E-5
²² Ne	1.0E-3	9.8E-4	9.1E-4	8.6E-4
²² Na	9.6E-5	6.9E-4	3.5E-4	1.9E-4
²³ Na	1.4E-3	1.6E-3	1.9E-3	1.9E-3
²⁴ Mg	2.0E-4	9.0E-4	4.1E-4	3.9E-4
²⁵ Mg	2.4E-3	4.6E-3	3.2E-3	1.3E-3
²⁶ Mg	2.8E-4	5.9E-4	5.1E-4	1.1E-3
²⁶ Al	5.4E-4	8.8E-4	6.5E-4	2.0E-4
²⁷ Al	2.0E-3	2.9E-3	2.2E-3	2.2E-3
²⁸ Si	5.6E-2	6.8E-2	5.8E-2	5.6E-2

Table 6. Main characteristics of the 1.35 M_⊙ ONe nova models compared with model ONe6 (José & Hernanz 1998)

	ONe6	ONe135A	ONe135B	ONe135C
Reaction Network	Old	A	B	C
ΔM_{env} (10^{-5} M _⊙)	0.54	0.54	0.54	0.54
T_{max} (10^8 K)	3.24	3.22	3.31	3.23
ΔM_{ejec} (10^{-5} M _⊙)	0.44	0.44	0.44	0.44
K (10^{45} erg)	0.9	1.0	1.0	1.0
²⁰ Ne	1.5E-1	1.3E-1	1.4E-1	1.4E-1
²¹ Ne	5.1E-5	9.9E-4	2.6E-4	3.9E-5
²² Ne	1.5E-4	1.9E-4	1.9E-4	1.7E-4
²² Na	6.0E-4	1.4E-3	1.0E-3	7.7E-4
²³ Na	6.6E-3	5.2E-3	5.5E-3	5.6E-3
²⁴ Mg	3.6E-4	8.6E-4	5.4E-4	5.4E-4
²⁵ Mg	4.2E-3	6.8E-3	4.5E-3	1.3E-3
²⁶ Mg	5.9E-4	6.3E-4	7.2E-4	2.1E-3
²⁶ Al	7.2E-4	1.1E-3	7.2E-4	1.5E-4
²⁷ Al	1.8E-3	2.1E-3	1.8E-3	2.0E-3
²⁸ Si	3.5E-2	5.0E-2	4.2E-2	4.1E-2

Fig. 1.— Snapshots of the evolution of $^{20,21,22}\text{Ne}$, $^{21,22,23}\text{Na}$, and $^{22,23}\text{Mg}$ (in mass fractions) along the accreted envelope for a $1.25\text{ M}_{\odot}$ ONe novae accreting at a rate $\dot{M} = 2 \times 10^{-10}\text{ M}_{\odot}\cdot\text{yr}^{-1}$. The mass coordinate represents the mass below the surface, relative to the total mass. The successive panels, from top to bottom, correspond to the time for which the temperature at the burning shell reaches 5×10^7 , 7×10^7 , 10^8 , 2×10^8 , $T_{\text{max}} = 2.44 \times 10^8$, plus 3 panels corresponding to the last phases of the explosion, when the white dwarf envelope has already expanded to a size of $R_{\text{wd}} \sim 10^9$, 10^{10} and 10^{12} cm, respectively. The arrow indicates the base of the ejected shells.

Fig. 2.— Main nuclear paths in both NeNa and MgAl cycles.

Fig. 3.— Same as Fig. 1, for the evolution of $^{24,25,26}\text{Mg}$, $^{26}\text{Al}^{g,m}$, $^{25,27}\text{Al}$, and $^{26,27}\text{Si}$.

Fig. 4.— Comparison between the nuclear reaction rate for $^{21}\text{Na}(p,\gamma)^{22}\text{Mg}$ provided in this work, with the analytic rate given in Caughlan & Fowler (1988).







